

WARNING

KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE



TERMIDOR[®] 100SC

Residual Insecticide

Contains 100 g/litre fipronil in the form of a suspension concentrate

GROUP 2 INSECTICIDE

For the control of ants, cockroaches, earwigs, houseflies, house crickets, web-spinning spiders, daddy-long-legs spiders, millipedes, slaters and termites (subterranean, damp wood and dry wood termites) in external areas and surrounds of domestic, commercial, public and industrial buildings and structures.

Contents: 1L

Registered to and distributed by:

BASF New Zealand Limited

NZBN 9429032151299

P.O. Box 407, Auckland 1140

Phone: 0800 932 273

crop-solutions.basf.co.nz

Hazard Classification

6.1D, 6.9B, 9.1A, 9.3C, 9.4A



WARNING

May be harmful if swallowed, inhaled or absorbed through the skin. May cause damage to the nervous system from repeated oral exposure at high doses. Very toxic to aquatic life with long lasting effects. Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates.

PRECAUTIONS

HANDLING: Take all reasonable steps to ensure that the product does not cause any significant adverse effects to the environment beyond the application area. This product must be under the personal control of a qualified pest control operator during use.

STORAGE: Store in original container, tightly closed, in a cool, well ventilated area away from food, feedstuffs, animals, or fertilisers, and under lock and key. DO NOT store product or diluted spray mix in direct sunlight and protect from frost. Stores containing more than 100 litres require emergency response plans, secondary containment systems, and are subject to signage.

TRANSPORT: DO NOT carry this product on a passenger service vehicle.

SPILLAGE: Collect spillage. Prevent any product from entering drains. Wear appropriate protective clothing. Absorb spill with an inert material (such as vermiculite, clay granules or synthetic absorbent) and place in a labelled, sealable container for disposal. Wash the spill area with water and detergent, then absorb the washings and place in the same sealable container. Dispose of the waste safely in a suitable landfill.

PERSONAL PROTECTION: When opening the container, preparing spray and using the spray mix, wear chemical resistant clothing, buttoned to the neck and wrist, a washable hat, boots, impervious gloves, and goggles or face shield.

GENERAL SAFETY AND HYGIENE MEASURES: Use only outdoors or in a well-ventilated area. Avoid contact with skin or inhalation of spray mist. Wash splashes of concentrate from skin and eyes immediately. DO NOT eat, drink or smoke while using this product. Wash the outside of gloves before removing. Remove protective clothing and wash hands and exposed skin thoroughly with soap and water before meals and after work. After each day's use, wash gloves, contaminated clothing, and face protection. Store work clothing separately.

USE AND ENVIRONMENTAL PROTECTION: Apply the spray with well-maintained and accurately calibrated equipment. DO NOT spray onto, over or into water. DO NOT allow spray drift outside the treatment area to occur. Take care to avoid runoff. DO NOT apply if heavy rains are expected to occur within 48 hours of application. Very toxic to bees. Spray must NOT contact plants in flower if they are likely to be visited by bees. Clean out application equipment thoroughly after use. DO NOT contaminate drains or waterways when disposing of equipment wash waters.

RE-ENTRY PERIOD: DO NOT enter treated areas until spray has dried.

RECORD KEEPING: Records of use must be kept.

NOTE: Storage, application and record keeping must be as described in the current version of the New Zealand Standard for the Management of Agrichemicals NZS8409.

DISPOSAL: Container: DO NOT reuse or refill the container. Triple rinse empty container and add rinsate to the spray tank. Recycle through Agrecovery (0800 247 326, www.agrecovery.co.nz). Product: DO NOT dispose of undiluted product on site. Dispose of this product only by using according to the label or at an approved landfill. DO NOT burn product or empty container. DO NOT contaminate water supply with chemical or empty container.

FIRST AID

If medical advice is needed, have product container or label at hand. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. IF SWALLOWED, DO NOT induce vomiting. Rinse mouth. IF ON SKIN or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. IF INHALED, remove to fresh air immediately and keep at rest in a position comfortable for breathing. IF ON EYES, wash out immediately with water. Get medical advice/attention if you feel unwell.

CONDITIONS OF SUPPLY

Read conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened. All goods supplied by BASF New Zealand Limited are of a high grade and conform to the chemical description of the label. However, to the extent permitted by law, neither BASF nor any of its Distributors accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with the supply of those goods including without limitation their storage, application, handling or use. BASF New Zealand Limited nor any of its Distributors, make no warranties of any kind, express or implied, with respect to fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter with respect to these goods. When you are a consumer as defined in the Consumer Guarantees Act 1993 (the "Act") these conditions will be subject to the provisions of the Act. However, the Act will not apply to goods supplied to you for business purposes.

Approved pursuant to the HSNO Act 1996, Code HSR100550
See www.epa.govt.nz for approval controls

MPI approved Type B (All animal product except dairy).

MPI approved for use in Farm Dairies and Recognized for use in Dairy Processing.

Batch No:

Date of Manufacture:

Registered to and distributed by:

BASF New Zealand Limited

NZBN 9429032151299

P.O. Box 407, Auckland 1140

Phone: 0800 932 273

crop-solutions.basf.co.nz

EMERGENCY PHONELINK
24 Hour Advice in an Emergency Only
0800 944 955



IIIIII BAR CODE IIIIII

® Registered Trademark of BASF
© Copyright 2019, BASF New Zealand Limited

GENERAL INFORMATION

TERMIDOR 100SC will control ants, cockroaches, web-spinning spiders, house flies and termites (subterranean, damp wood and dry wood termites) by direct contact and residual activity on treated surfaces. TERMIDOR 100SC is best applied as activity increases in early spring. Follow up applications during summer may be required.

RESISTANCE MANAGEMENT

TERMIDOR 100SC contains fipronil from the **GROUP 2 INSECTICIDE**. Some naturally-occurring insect biotypes resistant to TERMIDOR 100SC and other Group 2 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if TERMIDOR 100SC or other Group 2B insecticides are used repeatedly. The effectiveness of TERMIDOR 100SC on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, BASF New Zealand Ltd accepts no liability for any losses that may result from the failure of TERMIDOR 100SC to control resistant insects. TERMIDOR 100SC may be subject to specific resistance management strategies. For further information contact your BASF New Zealand Ltd representative.

DIRECTIONS FOR USE

Note: Before spraying advise residents to remove pets from the area to be treated. Cover or remove (as applicable) fish ponds, aquariums, bird cages, and open food and water containers before spraying. Doors and windows adjacent to the application site must be closed during surface application. DO NOT allow residents, children, other persons or pets into the immediate area during application. DO NOT allow people or pets to contact contaminated areas or to re-occupy contaminated areas until the clean-up is completed. An inspection should be carried out after treatment to ensure that no spillages or leaks have occurred and if any are identified these must be cleaned up before leaving the application site.

Restrains: Avoid runoff. DO NOT apply to excessively wet soils, immediately after or during heavy rain. DO NOT apply onto or around fruit trees and or edible crops. DO NOT apply at less than the label rate. DO NOT apply to internal/indoor surfaces. DO NOT apply onto or around fruit trees or edible plants.

Use Rate: Apply the prepared spray mix at either of the proposed use rates
6ml/L - 2 times / year
3ml/L - 4 times / year
6ml/L - 1 time / year + 3ml/L - 2 times / year

Pest	Situation	Critical Comments
<p>Nuisance Ants, including (but not limited to): Argentine Ant (<i>Linepithema humile</i>), Black House Ant (<i>Ochetellus glaber</i>), Pedicel Ant / Odorous House Ant (<i>Tapinoma</i> spp.), Pharaoh's Ant (<i>Monomorium phararonis</i>), Whitefooted Ant (<i>Technomyrmex albipes</i>).</p>	<p>External areas and surrounds of domestic, commercial, public and industrial buildings and structures</p>	<p>Treat surfaces 300 mm up and 300 mm out from where the building or structure touches the ground. Apply at the rate of 1 litre of spray mix per 25 lineal metres. Pay particular attention to potential entry points, such as weep holes, cracks and crevices. Also apply to ant trails and where ants are active away from the nest. Structures may include retaining walls, fences, garden borders, sheds etc.</p>
<p>Nesting Ants, including (but not limited to): Funnel Ant (<i>Aphaenogaster pythia</i>), Greenhead Ants (<i>Rhytidoponera</i> spp.), Meat Ants (<i>Iridomyrmex</i> spp.), Red Imported Fire Ant (<i>Solenopsis invicta</i>), Yellow Crazy Ant (<i>Anoplolepis gracilipes</i>).</p>	<p>Applications to nests.</p>	<p>Treat the nest entrance or mound, and where ants are active away from the nest. Apply at the rate 1 litre of spray mix per 16 m², or 60 ml per m².</p>
<p>Cockroaches, including (but not limited to): American Cockroaches (<i>Periplaneta americana</i>) Australian Cockroaches (<i>Periplaneta australasiae</i>) Oriental Cockroaches (<i>Blattella orientalis</i>) Smokybrown Cockroaches (<i>Periplaneta fuliginosa</i>)</p>	<p>External areas and surrounds of domestic, commercial, public and industrial buildings and structures</p>	<p>Treat external surfaces 300 mm up and 300 mm out from where the building or structure touches the ground. Apply at the rate of 1 litre spray mix per 25 lineal metres. Pay particular attention to potential entry points, such as weep holes, cracks and crevices. Also apply to where pest(s) may be active away from the building including retaining walls, fences, garden beds, sheds etc. Apply at the rate of 1 litre of spray mix per 16 m².</p>
<p>Web-spinning spiders, including (but not limited to): Black House Spider (<i>Baduma insignis</i>) and Red Back Spider (<i>Lacrodectus hasseltti</i>)</p>	<p>External areas and surrounds of domestic, commercial, public and industrial buildings and structures</p>	<p>Spray spiders, their webbing and areas where spiders may hide at the rate of 1 litre of spray mix per 16 m².</p>
<p>Daddy-Long-Legs Spiders (<i>Pholcus phalangioides</i>) Earwigs (<i>Forficula auricularia</i>) House Crickets (<i>Acheta domesticus</i>) Millipedes (<i>Julus hesperus</i>) Slaters (<i>Armadillidium vulgare</i>)</p>	<p>External areas and surrounds of domestic, commercial, public and industrial buildings and structures</p>	<p>Apply to the exterior of structures where pests enter, trail around the structure or where they crawl and hide. Apply around any exterior openings where pests could enter such as around doors, windows, vents, pipes, cracks, or drilled holes. May also be applied directly as a spot treatment.</p>

Pest	Situation	Critical Comments
Houseflies (<i>Musca domestica</i>)	External covered areas of domestic, commercial, public and industrial buildings and structures.	Apply to external surfaces where flies are likely to rest at the rate of 1 litre of spray mix per 16 m ² .
Termites (subterranean, damp wood and dry wood termites) including (but not limited to): <i>Coptotermes acinaciformis</i> , <i>Mastotermes darwiniensis</i> , <i>Schedorhinotermes</i> spp.	Pre-construction, Post-construction, Reticulation systems, protection of poles and fence posts, nests in poles and trees, wall cavity treatment.	Contact BASF for specific application advice.

MIXING

Half fill the spray tank with water and then add the required quantity of TERMIDOR 100SC. Stir and then top up the spray tank to the required volume. DO NOT use this product in a tank mix with other insecticides as the behaviour and efficacy of the product may be affected. Ensure equipment is free of leaks and clean from residues of other chemicals before mixing TERMIDOR 100SC. The use of a sprayer dedicated for the application of TERMIDOR 100SC is recommended.

IMPORTANT

Animal Product Processing Plants, Farm Dairies and Dairy Processing Plants: For use on external environment only. Not for use within processing facilities, near air inlets, water supplies, storage areas (including cleaning chemical storage) or packaging materials including pallets. Use in a manner that prevents the entry of the insecticide into edible product areas or stores containing packaging material.

® Registered Trademark of BASF
© Copyright 2019, BASF New Zealand Limited